

Engineering Mathematics I (Comp 400.001)
 Solution Set for Midterm Exam I : April 24, 2001

1. Find differential equations that describe the following curves.

- (a) (7 points) A curve in the xy -plane has the property that its slope at each point is equal to the sum of the square of the coordinates of the point.

$$y' = x^2 + y^2$$

- (b) (8 points) The graph of a nonnegative function has the property that the length of the arc between any two points on the graph is equal to the area of the region under the arc.

$$\int_a^x \sqrt{1+y'(t)^2} dt = \int_a^x y(t) dt$$

2. (10 points) Find the general solution of the following differential equation:

$$\frac{y^2}{2} + 2ye^x + (y + e^x) \frac{dy}{dx} = 0$$

$$\left(\underbrace{\frac{y^2}{2} + 2ye^x}_{\text{P}} \right) dx + \left(\underbrace{y + e^x}_{\text{Q}} \right) dy = 0 \quad (+2)$$

$$\frac{1}{Q} \left(\frac{\partial P}{\partial y} - \frac{\partial Q}{\partial x} \right) = \frac{1}{y+e^x} (y + 2e^x - e^x) = 1 \quad (+2)$$

$$F(x) = \exp \left(\int 1 dx \right) = e^x \quad (+2)$$

$$\left(\frac{1}{2}y^2 e^x + 2ye^x \right) dx + \left(ye^x + e^{2x} \right) dy = 0 \quad : \text{exact}$$

$$U(x, y) = \frac{1}{2}y^2 e^x + ye^{2x} + k_1(y) \quad (+2)$$

$$\frac{\partial U}{\partial y} = ye^x + e^{2x} + k_1'(y) = ye^x + e^{2x}$$

$$\therefore k_1(y) = \text{const}$$

$$\therefore U(x, y) = \frac{1}{2}y^2 e^x + ye^{2x} + c = 0 \quad (+2)$$

3. (10 points) Using a particular solution $y = e^{-x}$, find the general solution of the following differential equation:

$$xy'' + (x-1)y' - y = 0, \quad x > 0.$$

$$y'' + \underbrace{(\frac{1}{x})}_{P(x)} y' - \underbrace{\frac{1}{x}}_{q(x)} y = 0, \quad x > 0 \quad (+2)$$

$$U = \frac{1}{y_1^2} \cdot e^{-\int P(x) dx} = e^{2x} \cdot e^{-(x-\ln x)} = e^x \cdot x = xe^x \quad (+4)$$

$$y_2 = y_1 \int U(x) dx = e^{-x} \int xe^x dx = e^{-x} [xe^x - e^x] = (x-1) \quad (+4)$$

$$y = k_1 e^{-x} + k_2 (x-1)$$

4. (10 points) Find the general solution of the following equation:

$$y'' - 6y' + 9y = 6x^2 + 2 - 12e^{3x}.$$

$$\lambda^2 - 6\lambda + 9 = 0 \Rightarrow \lambda = 3 \quad](+2)$$

$$y_h = c_1 e^{3x} + c_2 x e^{3x}$$

$$y_p = a + bx + cx^2 + dx^2 e^{3x}$$

$$y'_p = b + 2cx + 2dx e^{3x} + 3dx^2 e^{3x}$$

$$y''_p = 2c + 2de^{3x} + 12dx^2 e^{3x} + 9dx^2 e^{3x}$$

$$y''_p - 6y'_p + 9y_p = 9a - 6b + 2c + (9b - 12c)x + 9cx^2 + 2de^{3x}$$

$$\left. \begin{array}{l} 9a - 6b + 2c = 2 \\ 9b - 12c = 0 \\ 9c = 6 \\ 2d = -12 \end{array} \right\} \Rightarrow \begin{array}{l} a = \frac{2}{3} \\ b = \frac{4}{9} \\ c = \frac{2}{3} \\ d = -6 \end{array}$$

$$\therefore y_p = \frac{2}{3} + \frac{4}{9}x + \frac{2}{3}x^2 - 6x^2 e^{3x} \quad (+5)$$

$$\therefore y = y_h + y_p = c_1 e^{3x} + c_2 x e^{3x} + \underline{\underline{\frac{2}{3} + \frac{4}{9}x + \frac{2}{3}x^2 - 6x^2 e^{3x}}}$$

5. (15 points) Find the general solution of the following equation:

$$x^2y'' - 3xy' + 3y = 2x^4e^x.$$

$$\begin{aligned} m(m-1) - 3m + 3 &= 0, \\ m = 1, 3 \quad] & \\ y_1 = x, \quad y_2 = x^3 & \end{aligned}$$

$$y_h = c_1x + c_2x^3$$

$$W = \begin{vmatrix} x & x^3 \\ 1 & 3x^2 \end{vmatrix} = 2x^3 \quad (+3)$$

$$y'' - \frac{3}{x}y' + \frac{3}{x^2}y = \underbrace{2x^2e^x}_{r(x)}$$

$$y_p = -y_1 \int \frac{y_2 - r(x)}{W(x)} dx + y_2 \int \frac{y_1 - r(x)}{W(x)} dx \quad (+3)$$

$$= -x \int \frac{x^3 \cdot 2x^2 e^x}{2x^3} dx + x^3 \int \frac{x \cdot 2x^2 e^x}{2x^3} dx$$

$$= -x \int x^2 e^x dx + x^3 \int x e^x dx$$

$$= -x (x^2 e^x - \int 2x e^x dx) + x^3 e^x$$

$$= x (2x e^x - \int 2 e^x dx)$$

$$= \underline{\underline{2x^2 e^x - 2x e^x}}.$$

$$\therefore y = \underline{\underline{c_1x + c_2x^3 + 2x^2 e^x - 2x e^x}} \quad (+2)$$

6. (10 points) Find the Laplace transform of the following function:

$$te^{2t}f'(t)$$

$$\begin{aligned}\mathcal{L}[f(t)] &= F(s) \\ \mathcal{L}[f'(t)] &= sF(s) - f(0) \quad \boxed{+3} \\ \mathcal{L}[e^{2t}f'(t)] &= (s-2)F(s-2) - f(0) \quad \boxed{+3} \\ \mathcal{L}[t(e^{2t}f'(t))] &= -\frac{d}{ds}[(s-2)F(s-2) - f(0)] \quad \boxed{+4} \\ &= -F(s-2) + (2-s)F'(s-2)\end{aligned}$$

7. (15 points) Find the Laplace transform of the following function:

$$t^2 \left(\int_0^t \tau \sin \tau d\tau \right)$$

$$\begin{aligned}\mathcal{L}[\sin t] &= \frac{1}{s^2 + 1} \quad \boxed{+2} \\ \mathcal{L}[t \sin t] &= -\frac{d}{ds} \left[\frac{1}{s^2 + 1} \right] = \frac{2s}{(s^2 + 1)^2} \quad \boxed{+2} \\ \mathcal{L}\left[\int_0^t \tau \sin \tau d\tau\right] &= \frac{1}{s} \cdot \frac{2s}{(s^2 + 1)^2} = \frac{2}{(s^2 + 1)^2} \quad \boxed{+4} \\ \mathcal{L}\left[t^2 \left(\int_0^t \tau \sin \tau d\tau \right)\right] &= \frac{d^2}{ds^2} \left[\frac{2}{(s^2 + 1)^2} \right] \quad \boxed{+4} \\ &= \frac{d}{ds} \left[\frac{-4(2s)}{(s^2 + 1)^3} \right] = \frac{-8(s^2 + 1)^3 + 8s \cdot 3(s^2 + 1)^2 \cdot 2s}{(s^2 + 1)^6} \quad \boxed{+3} \\ &= \frac{-8s^2 - 8 + 48s^2}{(s^2 + 1)^4} = \frac{40s^2 - 8}{(s^2 + 1)^4}\end{aligned}$$

8. (15 points) Find f and g satisfying the following simultaneous equations:

$$\begin{aligned}f(t) + \int_0^t (t-\tau)g(\tau)d\tau &= \sin 2t \\ g(t) + \int_0^t (t-\tau)f(\tau)d\tau &= 0\end{aligned}$$

$$\begin{aligned}\textcircled{+5} \quad \left\{ \begin{array}{lcl} F(s) + \frac{1}{s^2}G(s) & = & \frac{2}{s^2+4} \\ \frac{1}{s^2}F(s) + G(s) & = & 0 \end{array} \right. \Rightarrow \left\{ \begin{array}{lcl} s^2F(s) + G(s) & = & \frac{2s^2}{s^2+4} \\ F(s) + s^2G(s) & = & 0 \end{array} \right. \quad \textcircled{+3} \\ \left[\begin{array}{c} F(s) \\ G(s) \end{array} \right] = \frac{1}{s^4-1} \left[\begin{array}{cc} s^2 & -1 \\ -1 & s^2 \end{array} \right] \left[\begin{array}{c} \frac{2s^3}{s^2+4} \\ 0 \end{array} \right]\end{aligned}$$

$$\begin{aligned}F(s) &= \frac{2s^4}{(s^4-1)(s^2+4)} = \frac{1}{10} \cdot \frac{1}{s-1} + \frac{1}{10} \cdot \frac{1}{s+1} - \frac{1}{3} \cdot \frac{1}{s^2+1} + \frac{16}{15} \cdot \frac{2}{s^2+4} \quad \boxed{+4} \\ G(s) &= \frac{-2s^2}{(s^4-1)(s^2+4)} = -\frac{1}{10} \cdot \frac{1}{s-1} - \frac{1}{10} \cdot \frac{1}{s+1} - \frac{1}{3} \cdot \frac{1}{s^2+1} + \frac{4}{15} \cdot \frac{2}{s^2+4}\end{aligned}$$

$$\begin{aligned}f(t) &= \frac{1}{5} \cosh t - \frac{1}{3} \sin t + \frac{16}{15} \sin 2t \\ g(t) &= -\frac{1}{5} \cosh t - \frac{1}{3} \sin t + \frac{4}{15} \sin 2t \quad \boxed{+3}\end{aligned}$$